

# MaxiFlora

Commercial Fertilizer

0-0-6



**Guaranteed analysis:**

Soluble Potash (K<sub>2</sub>O) .....6.00%  
Sulfur (S) .....2.00%

**Derivation:**

Potash and Sulfur derived from Potassium Sulfate.

Weight per gallon = 8.80 Lbs.  
Kilograms per liter = 1.05 Kg.

**Handling:**

Shake or stir before use. Follow appropriate safety procedures. In case of accidental exposure, flush with plenty of water. Product is non-hazardous. Storage for more than one year is not recommended.

**Incompatibilities:**

Always jar test newly attempted mixes. No know incompatibilities.

**Directions for use:**

MaxiFlora is a supplemental nutrient for use with foliar spray applications. Properly timed applications may aid in improved crop nutrition. Use MaxiFlora with a soil fertility program and good cultural practices. The use of soil and tissue testing is recommended to determine the need for additional nutrients. Use 0.5 - 2.0 gallons per acre during the growing season, as needed.

**Directions for application:**

Application instructions: SHAKE WELL BEFORE USE. Apply with enough water to ensure thorough coverage.

For aircraft, use 5-10 gallons of water per acre and for ground applications use 10-250 gallons per acre. For best results apply MaxiFlora in the early mornings or late evening with a quality surfactant.

**General use guidelines:**

MaxiFlora may be applied to all flowering crops at 1-4 quarts per application and up to 4 applications per season. This includes, but is NOT limited to the following crops:

<u>Crop</u>	<u>Stage</u>	<u>Rate</u>	<u>Application*</u>
Apples, Pears	Bud Break	1 qt.	Do not apply to mature fruit
	First Green Leaf	1.5 qts.	
	3 week interval	1 qt.	
Apricot, Cherry, Peach, Plum, Nectarine	Bud Break	1.5 qts.	Do not apply to mature fruit
	First green leaf	2 qts.	
	3 week intervals	1.5 qts.	
Beans	Fourth true leaf	2 qts.	Make 4 appl.
	2 week intervals	1-2 qts.	
Broccoli, Cauliflower	Emergence	2 qts.	Directed spray Make 4 appl.
	2-3 week intervals	2 qts.	
Carrots	4-6 " Height	2 qts.	Make 4 appl.
	15-20 day interval	1-2 qts.	
Citrus	Pre bloom	2 qts.	Do not apply to mature fruit
	2 week interval	2 qts.	
Corn	6-16 " Height	2 qts.	
	2 week intervals	1.5 qts.	
Cotton	8-12 " Height	1-2 qts.	May be applied with pesticides
	2-3 week interval	1-2 qts.	
	Pre-cutout	2 qts.	
Cucumbers, Squash	3-4 leaf	1.1.5 qts.	Continue until 3 before harvest
	7-10 day interval	1 qt.	

Garlic	Leek stage	1.5 qts.	
	20-30 days later	1.5 qts.	
	15-20 days later	1.5 qts.	
Grapes	First leaf	1.5 qts.	
	Bud initiation	1 qt.	
	2-3 week interval	1-2 qts.	
Leafy vegetables Lettuce, Spinach	3-4 leaf	1-2 qts.	Apply after cutting
	2-3 week interval	1-2 qts.	
Melons	3 leaf	1-2 qts.	Continue until 3 weeks before harv.
	2-3 week interval	1-2 qts.	
Onions	3 leaf	2 qts.	
	2 week interval	2 qts.	
Potato, Radish	6-8 " height	1.5 qts.	
	2-3 week intervals	1.5 qts.	
Sorghum	Head elongation	1.5 qts.	
	2-3 week interval	1.5 qts.	
Sugarbeet	2-3 week interval	1-2 qts.	Make 4 appl.
Tomatoes	8-10 " Height	2 qts.	Make 4 appl.
	2 week interval	2 qts.	
Wheat, Barley	End of tillering beginning of stem elongation	2 qts.	

\* Always jar test newly attempted mixes.

**Notice to user:**

Call 1-800-USA-BIO1 for technical assistance concerning use rates for crops not mentioned in this label.

Manufactured by,  
**BioScientific, Inc.**  
Avondale, Arizona

5 GALLONS - 18.9 LITERS