

D & M Chem

Eclipse™ Calcium With Boron

Guaranteed Analysis

	By weight
Calcium(Ca)	25.00%
Boron(B)	0.10%

Derived from Calcium Carbonate and Boric Acid
 Contains 3.44 Lb. Ca., 0.014 Lb. B., per gallon
 Density: 13.75 Lbs. Per gallon @ 68 degrees F
 Do Not Store at Temperature Below 32° F.
 Sold in 250 gallon bulk shuttles
 Ph of 8.6

GENERAL INFORMATION

This is a non-phytotoxic micronutrient solution for foliar and soil application. The Calcium is highly available and absorbed by leaves and new, woody shoots, or the root system. Do not use aluminum or copper (including brass or bronze) components in any handling or storage equipment for this product. Mild steel components should not be used except for emergency service. Use stainless steel (316 or better), polyolefin (polyethylene, polypropylene, etc.), fiberglass, or other high-quality acid-resistant plastic materials. Check with equipment manufacturer for compatibility of components prior to use with this product.

General Recommendations

FOLIAR APPLICATIONS: For best absorption into the plant, apply this product in early morning or late evening. For aerial applications use at least 1 gallon of water for every pint of this product applied to the crop. Consult your local Farm advisor for information on sensitive varieties and hybrids.

D&M CHEM ECLIPSE is designed for use on crops grown on neutral to acid soils. D&M CHEM ECLIPSE is best used foliar or ground applied. Application of D&M CHEM ECLIPSE should begin prior to fruit formation and continue throughout harvest at 1 to 3 gallons per acre.

CAUTION: This fertilizer contains boron and is intended for use only on crops identified on this label. Its use on any other crops or under conditions other than those recommended may result in serious injury to crops.

Alfalfa	Carrots	Lettuce	Peas	Rice
Almonds	Celery	Melons	Peanuts	Safflower
Apples	Citrus	Milo	Pecans	Soybeans
Apricots	Corn	Mint	Peppers	Spinach
Asparagus	Cotton	Nectarines	Plums	Strawberries
Avocado	Cranberries	Nursery Stock	Pomegranates	Sugar Cane
Barley	Cucumbers	Oats	Potatoes	Sugar Beets
Beans (all)	Figs	Olives	Pasture &	Sweet Corn
Boysenberries	Garlic	Onions	Range	Tomatoes
Broccoli	Grapes	Ornamentals	Grass	Turf Grass
Brussels	Grass	Peaches	Prunes	Walnuts
sprouts	Hops	Pears	Rape	Watermelons
				Wheat

FRUIT AND NUT CROPS

APPLES AND PEARS, FILBERTS, PECANS, CITRUS, STONE FRUITS, GRAPES, AND OTHER FRUIT AND NUT CROPS: Foliage: Mix 1 to 3 gallons per acre with sufficient water and apply as a thorough cover spray. Applications should begin before bloom and continue through harvest at 1 to 3 gallons per acre.

FOR USE ON CROPS AND ORNAMENTALS

BEAN FOLAGE: Apply 1 to 3 gallons per acre prior to bloom on determinate and indeterminate varieties with post emergence herbicide application. Continue application at 1 to 3 gallons per acre until pods have formed.

COTTON FOLIAGE: Apply 1 to 3 gallons per acre prior to bloom with existing pesticide program (includes post emergence herbicides.) Continue application at 1 to 3 gallons per acre until first boll split.

POTATOES FOLIAR: Apply 1 to 3 gallons per acre prior to bloom. Continue application of 1 to 3 gallons per acre until 2 weeks prior to defoliation.

VEGETABLE CROPS: Apply 1 to 3 gallons per acre before fruit has formed. Continue applications of 1 to 3 gallons per acre until 2 weeks prior to harvest.

COLE CROPS: Same as Vegetable Crops.

OTHER CROPS AND USES

ORNAMENTALS, CONTAINER GROWN TREES AND SHRUBS, AND FIELD GROWN STOCK: Soil: Apply 4 to 16 ounces per 100 sq. ft. Repeat application at fertilization or regular intervals. Foliage: Apply 1 to 3 gallons per acre as a thorough cover spray.

GREENHOUSE-GROWN CHRYSANTHEMUMS, ROSES, CARNATIONS, AND POTTED FLORAL PLANTS: Soil Only: Apply 4 to 12 ounces per 100 sq. ft. at regular intervals with fertilizer or drip irrigation.

TURF GRASSES: Apply 1 gallon per acre or 4 to 8 oz. Per 1000 sq. ft. Mix with water, liquid fertilizer solution or pesticide solution and apply in early spring: Repeat as needed.

FOR APPLICATION IN IRRIGATION WATER

PIVOT SPRINKLER, OVERHEAD AND SOLID SET IRRIGATION: Apply 1 to 8 gallons per acre with irrigation water. The product may be incorporated into fertilizer and/or pesticide before injection. Use back-flow check valve to prevent material from running back into water system.

NOTICE

D&M Chem Inc. Warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of D&M Chem. In no case shall D&M Chem be liable for consequential, special or indirect damages resulting from the use of handling of this product. All such risks shall be assumed by the buyer. D&M Chem makes no warranties or Merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

MANUFACTURED BY
D&M CHEM INC.

112 Charron Road, Moxee, Washington 98936

"Information regarding the contents and levels of metals in this product is available on the Internet at: <http://www.aapfco.org/metals.htm>